

# THE LANCET

## **Supplementary appendix**

This appendix formed part of the original submission and has been peer reviewed. We post it as supplied by the authors.

Supplement to: Brose MS, Nutting CM, Jarzab B, et al, on behalf of the DECISION investigators. Sorafenib in radioactive iodine-refractory, locally advanced or metastatic differentiated thyroid cancer: a randomised, double-blind, phase 3 trial. *Lancet* 2014; published online April 24. [http://dx.doi.org/10.1016/S0140-6736\(14\)60421-9](http://dx.doi.org/10.1016/S0140-6736(14)60421-9).

# **Sorafenib in locally advanced or metastatic, radioactive iodine-refractory, differentiated thyroid cancer: a randomized, double-blind, phase 3 trial**

Marcia S Brose, Christopher M Nutting, Barbara Jarzab, Rossella Elisei, Salvatore Siena, Lars Bastholt, Christelle de la Fouchardiere, Furio Pacini, Ralf Paschke, Young Kee Shong, Steven I Sherman, Johannes WA Smit, John Chung, Christian Kappeler, Carol Pena, István Molnár, Martin J Schlumberger, on behalf of the DECISION Investigators\*

## **Supplementary Appendices**

### **Table of Contents**

- A. List of DECISION study investigators
- B. Randomization and dose modifications
- C. Primary and secondary endpoints
- D. Additional biomarker data

## **Appendix A. List of DECISION study investigators**

### **China**

Yi Wu, Fudan University Cancer Hospital

Zhu-Ming Guo, Sun Yat-Sen University Cancer Center

Fang Li, Peking Union Medical College Hospital

Yuankai Shi, Cancer Institute and Hospital of Chinese Academy of Medical

Ming Gao, Tianjin Medical University Cancer Institute and Hospital

Lin Li, West China Hospital of SiChuan University

ChunYan Zhang, Zhejiang Cancer Hospital

Jia-Dong Wang, Shanghai Renji Hospital

### **Japan**

Makoto Tahara, National Cancer Center Hospital East

Yasuhisa Hasegawa, Aichi Cancer Center Hospital

Shunji Takahashi, The Cancer Institute Hospital of JFCR

Yuichi Ando, Nagoya University Hospital

### **South Korea**

Young Kee Shong, Asan Medical Center

Jae Hoon Chung, Samsung Medical Center

Moo Il Kang, Seoul St. Mary's Hospital

Eun-Jig Lee, Severance Hospital Yonsei Univ College of Medicine

Do-Joon Park, Seoul National University Hospital

Min Ho Shong, Chungnam National University Hospital

### **Saudi Arabia**

Abdulrahman Al nuaim, King Faisal Specialist Hospital and Research Centre

### **Austria**

Markus Raderer, Medizinische Uni Wien

### **Belgium**

Ahmad Awada, Jules Bordet Institute

### **Bulgaria**

Tatiana Hadjieve, UMHAT "Tsaritsa Yoanna"

### **Denmark**

Lars Bastholt, Odense University Hospital

## **France**

Martin Schlumberger, Institut Gustave Roussy and University Paris Sud

Sophie Leboulleux, Institut Gustave Roussy and University Paris Sud

Francoise Bonichon, Institut Bergonie

Stephane Bardet, Centre François Baclesse

Christelle De la Fouchardiere, Consortium Cancer Thyroïdien, Hospices Civils-Centre Anticancéreux

Anthony Gonçalves, Institut Paoli Calmettes

Patrice Rodien, CHU Angers

Jean-Philippe Spano, Hôpital La Pitié Salpêtrière

Jean-Louis Wemeau, Hôpital Claude Huriez, CHRU Lille

## **Germany**

Daniela Schmidt, Universitätsklinikum Erlangen

Andreas Bockisch, Universitätsklinikum Essen- Nuklearmedizin

Martin Fassnacht, Universitätsklinikum Würzburg

Christine Spitzweg, Klinikum der Universität München

Oliver A. Cornely, Klinikum der Universität zu Köln

Ralf Paschke, Leipzig University

## **Italy**

Furio Pacini, University of Siena

Salvatore Siena, Ospedale Niguarda Ca' Granda

Rossella Elisei, University of Pisa

Efisio Puxeddu, Università degli studi di Perugia

Dario Giuffrida, Istituto Oncologico del Mediterraneo

Laura Fugazzola, Fondazione IRCCS Ospedale Maggiore

Anna Maria Colao, Università degli studi 'Federico II'

Lisa Licitra, Fondazione IRCCS Istituto Nazionale Tumori

Massimo Giusti, Università degli studi di Genova

## **The Netherlands**

Johannes Smit, Radboud University Medical Center

Thera Links, University Medical Center Groningen

Helena Kapiteijn, Leiden University Medical Center

## **Poland**

Jerzy Sowinski, Szpital Kliniczny im. Heliodora Swiecickiego Uniwersytetu M

Grzegorz Kaminski, Wojskowy Instytut Medyczny

Izabella Kozlowicz-Gudzinska, Centrum Onkologii Instytut im. Marii Skłodowskiej-Curie

Barbara Jarzab, Maria Skłodowska-Curie Memorial Cancer Centre and Institute of Oncology

### **Russia**

Viktor Medvedev, Medical Radiological Research Centre of RAMS

### **Spain**

Mariano Provencio, H.U.P Hierro-Majadahonda

Jaume Capdevila Castellón, H.U. Vall d'Hebron

### **Sweden**

Johanna Svensson, Sahlgrenska University Hospital

Jan Tennvall, Lunds Universitetssjukhus

Viveka Bergman, Linköpings Universitetssjukhus

Christel Hedman, Karolinska Univ Hospital

### **UK**

Christopher Nutting, Royal Marsden Hospital

Susan Clarke, Guy's Hospital

Georgina Gerrard, St James's University Hospital

Ujjwal Mallick, The Freeman Hospital-Northern Centre for Cancer Care

Laura Moss, Velindre Cancer Centre

Beng Yap, The Christie NHS Foundation Trust

Leslie Samuel, Aberdeen Royal Infirmary

Nicholas Reed, Beatson WoS Cancer Centre

### **US**

Douglas Adkins, Washington University Cancer Center

Omar Eton, Boston University Cancer Research Center

Marcia Brose, Abramson Cancer Center of the University of Pennsylvania

Steven Sherman, University of Texas MD - Anderson Cancer Center

Naifa Busaidy, University of Texas - MD. Anderson Cancer Center

Julie Sosa, Yale University School of Medicine Comprehensive Cancer Center

Renato Martins, University of Washington

A. Colevas, Stanford Cancer Center

Robert Ferris, University of Pittsburgh Center Institute

Francis Worden, University of Michigan Health System

Julie Bauman, University of New Mexico Cancer Center

Stephen Lim, Cedars-Sinai Medical Center

Taofeek Owonikoko, Emory Winship Cancer Institute

Krzysztof Misiukiewicz, Mount Sinai Medical Center

## **Appendix B. Randomization and dose modifications**

Randomization lists (one list for each of the six strata) were prepared by a separate Bayer randomization management (RM) group handling randomization tasks. All versions of the randomization list were stored electronically. The printout or the lists on the electronic storage media were stored by RM in a secure location with access only by the RM group, the external randomization process service provider, and the external provider responsible for providing data monitoring committee reviews. After primary completion of the study, the treatment information was released to Bayer data management for unblinding the study database.

**Table B1. Dose reduction levels for sorafenib and placebo**

	<b>Dose level 0</b>	<b>Dose level -1</b>	<b>Dose level -2</b>	<b>Dose level -3</b>
Sorafenib total daily dose, mg	800	600	400	200
Administration	2 x 200 mg tablets twice a day	2 x 200 mg and 1 x 200 mg tablet 12 hours apart (either could be given first)	1 x 200 mg tablet twice a day	1 x 200 mg tablet once a day
Placebo administration	2 tablets twice a day	2 tablets and 1 tablet 12 hours apart (either could be given first)	1 tablet twice a day	1 tablet once a day

**Table B2. Criteria for dose delay or dose modification of sorafenib or placebo due to haematologic adverse events**

Grade of haematologic adverse event	Dose delay	Dose modification
Grade 0–2	No delay	No change
Grade 3	No delay	DECREASED one dose level <sup>b</sup>
Grade 4	DELAYED until ≤grade 2 <sup>a</sup>	DECREASED two dose levels <sup>b</sup>

<sup>a</sup>If no recovery after 30-day delay, treatment was discontinued unless the patient was deriving clinical benefit.

<sup>b</sup>If another dose reduction after dose level -3 was required, treatment was discontinued.

**Table B3. Criteria for dose delay or dose modification of sorafenib or placebo due to nonhaematologic adverse events (except skin toxicity and hypertension)<sup>a</sup>**

Grade of nonhaematologic adverse event <sup>a</sup>	Dose delay	Dose modification
Grade 0–1	No delay	No change
Grade 2	No delay	DECREASED one dose level <sup>c,d</sup>
Grade 3: 1 <sup>st</sup> occurrence	DELAYED <sup>b</sup> until ≤grade 2	DECREASED one dose level <sup>c,d</sup>
Grade 3: no improvement within 7 days, or 2 <sup>nd</sup> or 3 <sup>rd</sup> occurrence	DELAYED <sup>b</sup> until ≤grade 2	DECREASED two dose levels <sup>c,d</sup>
Grade 3: 4 <sup>th</sup> occurrence		DECREASED three dose levels <sup>c,d</sup>
Grade 4	Discontinued from protocol therapy	Discontinued from protocol therapy

<sup>a</sup>Also excluded nausea/vomiting that had not been premedicated, and diarrhoea.

<sup>b</sup>If no recovery after 30-day delay, treatment was discontinued unless patient was deriving clinical benefit.

<sup>c</sup>If another dose reduction after dose level -3 was required, treatment was discontinued.

<sup>d</sup>For patients who required a dose reduction for grade 2 or grade 3 toxicities, the dose of study drug may have been increased to the starting dose or up one dose level after one full cycle of therapy had been administered with the reduced dose without the appearance of the toxicity >grade 1.

**Table B4. Criteria for dose modification of sorafenib or placebo due to hypertension**

CTCAE grade of hypertension	Management/next dose
Grade 1	Increased BP monitoring considered
Grade 2 asymptomatic and diastolic BP <110 mm Hg	Begin antihypertensive therapy and continue study drug
Grade 2 symptomatic/persistent or diastolic BP $\geq$ 110 mm Hg or Grade 3	1. Study drug delayed <sup>a</sup> until symptoms resolved and diastolic BP $\leq$ 100 mm Hg, and patient treated with antihypertensives. When the study drug was restarted, it was reduced by one dose level <sup>b</sup> 2. If diastolic BP not controlled ( $\leq$ 100 mm Hg) on therapy, study drug was reduced another dose level <sup>c</sup>
Grade 4	Discontinued from protocol therapy

BP, blood pressure; CTCAE, Common Terminology Criteria for Adverse Events

<sup>a</sup>Patients requiring a delay of >14 days had to discontinue study drug.

<sup>b</sup>Patients may have been able to resume full dose later once BP was adequately controlled.

<sup>c</sup>Patients requiring >2 dose reductions had to discontinue study drug.

**Table B5. Criteria for dose modification of sorafenib or placebo due to skin toxicity**

<b>Grade of skin toxicity<sup>a</sup></b>		<b>Suggested dose modification</b>
Grade 1	Any occurrence	Maintained dose level and instituted supportive measures immediately for symptomatic relief
Grade 2 <sup>b</sup>	1 <sup>st</sup> occurrence	Instituted supportive measures immediately and considered a decrease of sorafenib dose by one dose level. If no improvement within 7 days, see below
	No improvement within 7 days or 2 <sup>nd</sup> occurrence	Interrupted until resolved to grade 0–1 When treatment resumed, decreased dose by one dose level
	3 <sup>rd</sup> occurrence	Interrupted until resolved to grade 0–1 When treatment resumed, decreased dose by two dose levels
	4 <sup>th</sup> occurrence	Discontinued from protocol therapy
Grade 3 <sup>b</sup>	1 <sup>st</sup> occurrence	Interrupted until resolved to grade 0–1 When treatment resumed, decreased dose by one dose level
	2 <sup>nd</sup> occurrence	Interrupted until resolved to grade 0–1 When treatment resumed, decreased dose by two dose levels
	3 <sup>rd</sup> occurrence	Discontinued from protocol therapy

<sup>a</sup>Dermatologic events were graded according to the Common Terminology Criteria for Adverse Events with the exception of hand–foot skin reactions, which were graded as follows:

grade 1 (numbness, dysesthesia/paresthesia, tingling, painless swelling or erythema of the hands and/or feet and/or discomfort that did not disrupt normal activities);

grade 2 (painful erythema and swelling of the hands and/or feet and/or discomfort that affected the patient's activities);

grade 3 (moist desquamation, ulceration, blistering or severe pain of the hands and/or feet and/or severe discomfort that caused the patient to be unable to work or perform activities of daily living).

<sup>b</sup>For patients who required a dose reduction for grade 2 or 3 rash or HFSR, the dose of study drug may have been increased to the starting dose after one full cycle of reduced dose therapy had been administered and there had been no appearance of rash or HFSR  $\geq$  grade 1.

## Appendix C: Primary and key secondary endpoints

The primary endpoint was progression-free survival, assessed every 8 weeks by central independent blinded review using Response Evaluation Criteria in Solid Tumors (RECIST) v1.0, from the date of randomization to the date of radiological progression or death. Radiological progression in bone as defined in RECIST v1.0 was modified specifically in this protocol as follows: 1) radiological appearance of new lesions; 2)  $\geq 20\%$  increase in the sum of the longest diameter of all target lesions, which may include bone lesions if they have measurable soft tissue components; 3) bone lesions that require external radiation. Secondary endpoints included overall survival (measured from the date of randomization to the date of death), time to progression (TTP; measured from the date of randomization to the date of radiological progression), objective response rate (complete or partial response), disease control rate (complete or partial response, or stable disease), and duration of response (defined as the time from the first documented objective response until disease progression or death). FDG-PET scan at baseline was required in centres with access to PET scanners. PFS, OS and TTP were analyzed in all randomized patients. ORR and DCR were analyzed in patients who received study medication and had a baseline and a post-baseline tumor evaluation.

Tumour response and progression-free survival were evaluated by both central review and investigator assessment. Primary and secondary efficacy endpoints were based on central review during the double-blind period; unblinding and starting open-label sorafenib was decided by investigator assessment.

## 1 Appendix D. Additional biomarker data

### 2 Table D1. Mutations tested in the HRAS, KRAS, NRAS, and BRAF genes

Oncogene	Mutations assayed
<i>HRAS</i>	G12V/D, G13C/R/S, Q61H/H, Q61L/R/P, Q61K
<i>KRAS</i>	G12C, G12R, G12S, G12V, G12A, G12F, G13V/D, A59T, Q61E/K, Q61L/R/P, Q61H/H
<i>NRAS</i>	G12V/A/D, G12C/R/S, G13V/A/D, G13C/R/S, A18T, Q61L/R/P, Q61H, Q61E/K
<i>BRAF</i>	G464R, G464V/E, G466R, F468C, G469S, G469E, G469A, G469V, G469R, D594V/G, F595L, G596R, L597S, L597R, L597Q, T599I, V600E, V600K, V600L, K601N, K601E

3

4 **Table D2. Demographic and clinical characteristics of the subpopulation for genetic analysis, compared with the overall study**  
5 **population**

	Overall population (N=417)	Subpopulation for genetic analysis		
		Overall (n=256)	Sorafenib (n=126)	Placebo (n=130)
Sorafenib PFS benefit				
Hazard ratio	0.59	0.57		
95% confidence interval	0.45–0.76	0.42–0.78	NA	NA
P value	<0.001	<0.001		
Female, n (%)	218 (52.3)	129 (50.4)	59 (46.8)	70 (53.8)
Age, median (range)	63 (24–87)	63 (24–87)	63 (24–81)	64 (30–87)
Ethnicity, n (%)				
White	251 (60.2)	180 (70.3)	86 (68.3)	94 (72.3)
Asian	99 (23.7)	29 (11.3)	14 (11.1)	15 (11.5)
Black	11 (2.6)	9 (3.5)	5 (4.0)	4 (3.1)
Other or not reported	56 (13.4)	38 (14.8)	21 (16.7)	17 (13.1)
ECOG performance status, n (%)				
0	259 (62.1)	162 (63.3)	80 (63.5)	82 (63.1)
1	143 (34.3)	85 (33.2)	41 (32.5)	44 (33.8)
2	13 (3.1)	7 (2.7)	4 (3.2)	3 (2.3)
Histology by central review, n (%)				
Papillary	237 (56.8)	156 (60.9)	74 (58.7)	82 (63.1)
Follicular	106 (25.4)	64 (25.0)	31 (24.6)	33 (25.4)
Poorly differentiated	40 (9.6)	32 (12.5)	18 (14.3)	14 (10.8)
Well differentiated	3 (0.7)	1 (0.4)	1 (0.8)	0
Nonthyroid	1 (0.2)	0	0	0
Medullary	1 (0.2)	0	0	0
Oncocytic carcinoma	2 (0.5)	1 (0.4)	1 (0.8)	0
Carcinoma, not otherwise specified	3 (0.7)	0	0	0
Missing/nondiagnostic	27 (6.5)	2 (0.8)	1 (0.8)	1 (0.8)

6 ECOG, Eastern Cooperative Oncology Group; NA, not applicable

7 **Table D3. Prognostic significance of BRAF and RAS mutation status on progression-free survival: multivariate model in**  
8 **patients treated with sorafenib and placebo including BRAF, RAS, clinical variables, and histology as covariates**

Variable	Level	Progression-free survival					
		Full analysis			Papillary patients only		
		HR	95% CI	P-value*	HR	95% CI	P-value*
N (events)		254 (154)			155 (89)		
Treatment	Sorafenib/placebo	0.50	0.36–0.69	<0.001	0.46	0.30–0.72	<0.001
Sex	Male/female	1.10	0.80–1.52	0.559	1.05	0.68–1.61	0.825
Ethnic origin	Other/white	0.76	0.48–1.21	0.246	0.77	0.43–1.38	0.379
	Not reported/white	0.86	0.54–1.38	0.539	0.85	0.45–1.61	0.620
Age		0.99	0.97–1.00	0.048	0.98	0.96–1.00	0.068
Histology	Follicular/papillary	1.37	0.90–2.07	0.138			
	Poorly differentiated/papillary	1.69	1.03–2.77	0.039			
	Other/papillary	1.02	0.31–3.39	0.972			
ECOG PS	1+2/0	1.36	0.97–1.92	0.075	0.92	0.58–1.45	0.714
BRAF	Mutation/wild-type	0.70	0.45–1.09	0.119	0.67	0.41–1.10	0.115
RAS	Mutation/wild-type	1.38	0.90–2.11	0.135	1.69	0.93–3.05	0.083


9 CI, confidence interval; ECOG PS, Eastern Cooperative Oncology Group performance status; HR, hazard ratio

10 \* P-values uncorrected for multiple hypothesis testing.

**Figure D1: Predictive analysis of biomarkers**


Kaplan–Meier graphs of progression-free survival by biomarker subgroups: BRAF mutation (P-value for biomarker-treatment interaction =0.653) [panels a and b]; RAS mutation (P-value for biomarker-treatment interaction =0.422) [panels c and d]; and thyroglobulin P-value for biomarker-treatment interaction =0.909) [panels e and f]. Similar results were seen when thyroglobulin was analysed as a continuous variable (P-value for biomarker-treatment interaction =0.988). Baseline median thyroglobulin =449.4 ng/ml.

a) BRAF wild-type




BRAF	Treatment	n	Events	Median PFS, days (months)	95% CI
WT	Sorafenib	92	59	271 (8.9)	170-337
WT	Placebo	87	64	117 (3.8)	106-175

b) BRAF mutation




BRAF	Treatment	n	Events	Median PFS, days (months)	95% CI
MT	Sorafenib	34	14	623 (20.5)	393-NR
MT	Placebo	43	25	286 (9.4)	174-NR

c) RAS wild-type




RAS	Treatment	n	Events	Median PFS, days (months)	95% CI
WT	Sorafenib	102	60	329 (10.8)	272-451
WT	Placebo	104	69	175 (5.7)	113-284

d) RAS mutation




RAS	Treatment	n	Events	Median PFS, days (months)	95% CI
MT	Sorafenib	24	13	167 (5.5)	146-NR
MT	Placebo	26	20	105 (3.4)	69-175

e) Low baseline TG



BL TG	Treatment	n	Events	Median PFS, days (months)	95% CI
Low	Sorafenib	98	60	556 (18.3)	354-722
Low	Placebo	103	60	268 (8.8)	223-391

f) High baseline TG



BL TG	Treatment	n	Events	Median PFS, days (months)	95% CI
High	Sorafenib	106	68	221 (7.3)	168-294
High	Placebo	96	70	113 (3.7)	105-169